

Secondary Publication



Clemens, Vera; Berthold, Oliver; Witt, Andreas; u. a.

Lifespan risks of growing up in a family with mental illness or substance abuse

Date of secondary publication: 03.06.2026

Version of Record (Published Version), Article

Persistent identifier: urn:nbn:de:bvb:473-irb-115396x

Primary publication

Clemens, Vera; Berthold, Oliver; Witt, Andreas; u. a. (2020): Lifespan risks of growing up in a family with mental illness or substance abuse, in: Scientific reports, London: Springer Nature, Vol. 10, No. 15453, pp. 1–8, doi: 10.1038/s41598-020-72064-w.

Legal Notice

This work is protected by copyright and/or the indication of a licence. You are free to use this work in any way permitted by the copyright and/or the licence that applies to your usage. For other uses, you must obtain permission from the rights-holders.

This document is made available under a Creative Commons license.



The license information is available online:

<https://creativecommons.org/licenses/by/4.0/legalcode>



OPEN

Lifespan risks of growing up in a family with mental illness or substance abuse

Vera Clemens¹✉, Oliver Berthold¹, Andreas Witt¹, Cedric Sachser¹, Elmar Brähler^{2,3}, Paul L. Plener^{1,4}, Bernhard Strauß⁵ & Jörg M. Fegert¹

Growing up in a family with one member being affected by mental health problems or substance abuse is an adverse childhood experience which can lead to socioeconomic and health-related impairments in later life. Furthermore, the risk of child maltreatment is increased in affected families, which often adds to the individual risk factors. However, the interdependence between the particular risk factors is not well understood. To examine the correlation between mental health problems or substance abuse and child maltreatment within families and long term consequences for affected children, a cross sectional population representative survey in Germany (N = 2,531) has been conducted. The risk of child maltreatment was 5 to 5.6 times higher if mental illness and 4.9 to 6.9 times higher if substance abuse of a family member was reported. Furthermore, the risk of health problems, including obesity, decreased life satisfaction, lower income, low educational achievement, unemployment and living without a partner was increased if participants grew up in a family affected by mental health problems or substance abuse. All associations were mediated significantly by child maltreatment. These results point towards an urgent need for greater awareness for child protection issues in families affected by mental health problems or substance abuse.

Adverse childhood experiences (ACE) can affect life in different ways. The long-term consequences are widespread and can include psychosocial and economic impairments, a significant reduction in quality of life and increased morbidity due to both, mental and somatic health problems¹. In his famous ACE study 20 years ago, Vincent Felitti demonstrated that with increasing number of ACEs the risk of suicide attempts, smoking, alcohol and drug consumption and many diseases increased up to 12-fold². Recently published studies on population representative samples in Germany and the U.S. show that child maltreatment is associated with an increased risk for leading causes of morbidity and mortality, including cardiovascular and oncological diseases^{3,4}. Overall, ACEs can lead to a shortening of the life span of up to 20 years⁵. In addition to these serious consequences for the life of each individual affected, they lead to an enormous economic burden with annual expenditures between 11 and 30 billion euros in Germany alone⁶. Taken together, ACEs can be considered as major public health problems.

ACEs include all forms of child maltreatment, defined as any intentional acts of commission or omission that cause harm or expose the child/adolescent to the risk of harm—regardless of the intention to harm⁷. Child maltreatment is frequent—a recent epidemiologic study from a German sample reports at least moderate experience of emotional abuse in 6.6%, physical abuse in 6.7%, sexual abuse in 7.6% and emotional and physical neglect in 13.3% and 22.5%, respectively⁸. The term ACE also includes stressful childhood experiences such as intimate partner violence between parents, parental separation or divorce, and being part of a family affected by mental health problems or substance abuse. The last two items are of particular relevance to mental health professionals, as 10–30% of inpatients with psychiatric disorders have underage children^{9,10}. A better understanding of the possible long-term consequences of growing up in families affected by mental health problems or substance abuse is essential for effective prevention.

¹Department of Child and Adolescent Psychiatry/Psychotherapy, University of Ulm, Steinhövelstr. 5, 89073 Ulm, Germany. ²Department for Psychosomatic Medicine and Psychotherapy, University Medical Center of Johannes Gutenberg University of Mainz, Mainz, Germany. ³Department of Medical Psychology and Medical Sociology, University of Leipzig, Leipzig, Germany. ⁴Department for Child and Adolescent Psychiatry, Medical University of Vienna, Vienna, Austria. ⁵Institute of Psychosocial Medicine and Psychotherapy, Jena University Hospital, Jena, Germany. ✉email: vera.clemens@uni-ulm.de

Earlier research showed that families affected by mental health problems or substance abuse are at higher risk for child maltreatment^{11–13}. As shown above, child maltreatment is a powerful risk factor on its own^{1,3,14,15}. For effective prevention, we need to better understand the interplay between these risk factors, namely growing up in a family affected by mental health problems and substance abuse, and child maltreatment. In a previous analysis, we were able to show that child maltreatment mediates the risk of household dysfunction for depression, anxiety, life quality and overall health status¹⁶. Based on these associations, we hypothesized that child maltreatment partly mediates the long term consequences of growing up in a family affected by mental health problems and substance abuse.

Therefore, we examined health and socio-economic indicators in adulthood as well as the experience of child maltreatment in dependence of growing up in a family affected by mental health problems or substance abuse in a sample of the German population representative in regard of gender and age. A mediation analysis for child maltreatment in relation to growing up in a family affected by mental health problems or substance abuse.

Methods

A representative sample of the German population was randomly collected by a social research institute (USUMA, Berlin, Germany) as described elsewhere¹⁷.

Data collection took place between November 2017 and February 2018. The sample was representative of the German population over 14 years of age in terms of age and gender. Out of 5,160 identified addresses, 2,531 persons participated in the study (response rate: 49.1%).

All participants received information and written consent. In the case of minors, the participants gave their consent after clarification with the consent of their caregivers. The study was conducted in accordance with the Declaration of Helsinki and complied with the ethical guidelines of the International Code of Marketing and Social Research Practice of the International Chamber of Commerce and the European Society of Opinion and Marketing Research. The study was approved by the Ethics Commission of the Medical Faculty of the University of Leipzig.

Measures. The prevalence of different forms of ACEs was assessed using the German version of the Adverse Childhood Experiences Questionnaire, a standard tool for retrospective assessment of ACEs with satisfactory internal consistency (Cronbachs $\alpha = 0.76$)¹⁸. The questions relating growing up in a family affected by mental health problems or substance abuse were used as independent variables in all analyses. The wording of the questions was: "Did you live with someone who had alcohol problems, was addicted to alcohol or drugs?" and "Did a member of your household suffer from depression, mental health problems or did attempt suicide?"

Different psychosocial and health-related variables served as outcomes. In detail, satisfaction with life was assessed based on the question "How satisfied are you at present, all in all, with your life", scale 1 (not satisfied at all) to 11 (completely satisfied) according to Beierlein and colleagues¹⁹. The state of health was assessed using the EuroQOL 5-Dimensions 3-Level (EQ-5D-3L) questionnaire. The EQ-5D-3L covers five dimensions on current problems: "Mobility", "Self-care", "Usual Activities", "Pain/Physical discomfort" and "Anxiety/Depression". The answers encompass 3 levels: no problems, moderate problems, or severe problems²⁰. For the EQ-5D-3L index, all 5 dimensions can be summarized using a formula based on country-specific weights for the individual items²¹. In the present study, weights were used which were derived from a large representative German population sample²². Only valid answers were included into the respective analysis. The number of included cases is given for each analysis. The participants were on average 48.6 years old ($SD = 18.0$), 56.4% were women. 96% stated that they had German citizenship. The sample was representative of the German population over the age of 14 in terms of age and gender. Sample characteristics are presented in Table 1.

Statistical analyses. All statistical analyses were performed with SPSS version 21. *p* values correspond to two-tailed tests. The prevalence rates were determined using descriptive analyses. Binary logistic regression analyses were performed in order to analyze the odds for the assessed long-term outcomes if mental illness or substance abuse were present in the family of origin. Age, gender and educational achievement were included into the analyses as co-variates. If school qualification served as outcome of regression analysis, no educational achievement was not included as confounder. Bonferroni corrections for multiple comparisons were performed to guard against type I error inflation and α -levels were adjusted accordingly.

The mediation analyses were carried out with the Hayes PROCESS macro²³. Path analyses were performed with 5,000 bootstrapping samples. The number of different ACE subtypes (0–5) served as a mediation variable. As potential confounders, age, gender and educational achievement were included into the analyses.

Results

In total, 267 (10.5%) and 421 (16.6%) of participants reported growing up in a family affected by mental health problems, or substance abuse, respectively. 316 (12.5%) of the participants reported emotional and 230 (9.1%) physical abuse, 109 (4.3%) sexual abuse, 338 (13.4%) emotional and 109 (4.3%) physical neglect (see Table 1).

Child maltreatment is more frequent in families affected by mental health problems or substance abuse. There was a positive association between experiences of maltreatment and growing up in a family affected by mental health problems or substance abuse. Mental health problems in the family was associated with an increased risk for all subtypes of child maltreatment (ORs 5.07–5.63), as was substance abuse (ORs 4.87–6.91). For growing up in a family affected by substance abuse, the risk of physical abuse (OR 6.81) and physical neglect (OR 6.91) increased the most. If growing up in a family affected by mental health problems was reported, the risk of emotional abuse (OR 5.63) was highest (see Table 2).

	Total (N = 2,531)
Age (M, SD)	48.6 (18.0)
Gender (n, %)	
Female	1,401 (55.4)
Male	1,130 (44.6)
German citizenship (n, %)	2,429 (96.0)
Living with a partner (n, %)	1,351 (53.4)
Number of periods of unemployment (lifetime) (M, SD)	0.79 (1.51)
Monthly household income in € (M, SD)	1759.56 (715.25)
Life satisfaction (M, SD)	8.26 (2.07)
EQ-5D-3L index (M, SD)	0.92 (0.14)
Highest level of education (n, %)	
Left school before graduation	56 (2.3)
School graduation	2,186 (75.8)
German baccalaureate	307 (12.5)
University degree	233 (9.5)
Adverse childhood experiences (ACEs) (n, %)	
Emotional abuse	316 (12.5)
Physical abuse	230 (9.1)
Sexual abuse	109 (4.3)
Emotional neglect	338 (13.4)
Physical neglect	109 (4.3)
Substance abuse of a family member	421 (16.6)
Mental illness of a family member	267 (10.5)

Table 1. Sample characteristics. Presented as mean (M) ± standard deviation (SD) or number of participants (%).

	Emotional abuse	Physical abuse	Sexual abuse	Emotional neglect	Physical neglect
Substance abuse					
OR (95% CI)	5.91 (4.55;7.69)	6.81 (5.07;9.13)	4.87 (3.26;7.27)	5.06 (3.92;6.53)	6.91 (4.61;10.35)
Wald	176.86	162.44	59.89	155.34	87.64
Nagelkerkes R ²	0.14	0.16	0.12	0.12	0.14
<i>p</i> value	<0.001	<0.001	<0.001	<0.001	<0.001
N	2,445	2,441	2,442	2,442	2,443
Mental health problems					
OR (95% CI)	5.63 (4.20;7.54)	5.07 (3.67;7.01)	5.27 (3.44;8.08)	5.13 (3.84;6.84)	5.22 (3.40;8.00)
Wald	133.54	96.69	58.07	123.32	57.09
Nagelkerkes R ²	0.11	0.10	0.11	0.10	0.09
<i>p</i> value	<0.001	<0.001	<0.001	<0.001	<0.001
N	2,432	2,428	2,430	2,431	2,431

Table 2. Frequency and risk of child abuse depending on growing up in a family affected by mental health problems or substance abuse. Presented as OR (odds ratio) adjusted for gender, age and educational achievement. Bonferroni correction was performed, *p* < 0.01 was considered as statistically significant. 95% CI = 95% confidence interval.

Higher risk of poor health and low socioeconomic status in families affected by mental health problems illness or substance abuse. There was a positive correlation between growing up in a family affected by mental health problems or substance abuse and poorer health, obesity and lower socioeconomic status.

Participants were more than twice as likely to be dissatisfied with their lives if they grew up in a family affected by mental health problems or substance abuse. Furthermore there was a positive correlation with not living with a partner, having been unemployed at least once in a lifetime and lower income. Only growing up in a family affected by substance abuse was positively related with low educational achievement (see Table 3).

	Poor Health (< 25. Percentile.)	Low life satisfaction (≤ 7)	Low educational achievement	Not living with a partner	At least once unemployed	Monthly equalized income below 1,250 €	Obesity
Substance abuse							
OR (95% CI)	2.58 (1.93;3.46)	2.26 (1.76;2.91)	2.50 (1.42;4.42)	1.45 (1.17;1.79)	2.34 (1.86;2.95)	1.83 (1.46;2.30)	1.49 (1.12;1.97)
<i>p</i> value	<0.001	<0.001	0.002	0.001	<0.001	<0.001	0.006
Nagelkerkes R ²	0.25	0.05	0.03	0.01	0.07	0.07	0.03
N	2,375	2,390	2,453	2,439	2,298	2,387	2,505
Mental health problems							
OR (95% CI)	2.66 (2.08;3.41)	2.82 (2.12;3.76)	1.60 (0.77;3.30)	1.57 (1.21;2.04)	1.58 (1.21;2.07)	1.92 (1.46;2.54)	1.42 (1.01;1.98)
<i>p</i> value	<0.001	<0.001	0.207 (n.s.)	0.001	0.001	<0.001	0.042 (n.s.)
Nagelkerkes R ²	0.26	0.05	0.02	0.01	0.05	0.06	0.03
N	2,364	2,377	2,440	2,426	2,287	2,374	2,492

Table 3. Frequency and risk of health and socio-economic impairments depending on living with household members who were substance abusers or mentally ill before the age of 18, adjusted for gender, age and educational achievement. Bonferroni correction was performed, $p < 0.007$ was considered as statistically significant. Presented as OR (odds ratio). 95% CI = 95% confidence interval.

	Total effect	Direct effect	Indirect effect (mediation)	% of total effect is mediated
Mental illness				
Health (N = 2,344)	-0.05 (-0.07; -0.03)	-0.03 (-0.05; -0.01)	-0.02 (-0.03; -0.0)	40
Life satisfaction (N = 2,350)	-1.16 (-1.42; -0.90)	-0.77 (-1.05; -0.50)	-0.39 (-0.52; -0.28)	34
Income (N = 2,348)	-191.30 (-281.93; -100.66)	-120.35 (-214.39; -25.68)	-71.26 (-101.18; -43.98)	37
Unemployment (N = 2,411)	0.31 (0.12;0.51)	0.11 (-0.10;0.31)	0.11 (0.13;0.29)	35
BMI (N = 2,394)	n.s.	n.s.	0.31 (0.04;0.56)	-
Substance abuse				
Health (N = 2,335)	-0.05 (-0.07; -0.03)	-0.03 (-0.05; -0.02)	-0.02 (-0.03; -0.01)	40
Life satisfaction (N = 2,361)	-0.93 (-1.15; -0.71)	-0.55 (-0.78; -0.32)	-0.38 (-0.50; -0.27)	41
Income (N = 2,359)	-214.20 (-288.03; -140.37)	-152.81 (-231.66; -73.95)	-61.39 (-88.53; -34.64)	29
Unemployment (N = 2,422)	0.52 (0.36;0.68)	0.36 (0.19;0.53)	0.16 (0.09;0.24)	31
BMI (N = 2,406)	n.s.	n.s.	0.32 (0.10;0.55)	-

Table 4. Association between growing up in a family affected by mental health problems or substance abuse, child abuse and impairments in adulthood. Mediation analyses of associations between growing up in a family affected by mental health problems or substance abuse and general health (via EQ-5D-3L index), life satisfaction, normalized income and the frequency of unemployment. Total effect = association between growing up in a family affected by mental health problems or substance abuse and a particular outcome variable without exclusion of child maltreatment. Direct effect = association between growing up in a family affected by mental health problems or substance abuse and a particular outcome variable after exclusion of child maltreatment. Indirect effect = association between the number of experienced forms of child maltreatment with the respective outcome variable. Presented as β -coefficient and 95% confidence interval, adjusted for gender, age and educational achievement. Bonferroni correction was performed, $p < 0.01$ was considered as statistically significant. Results are significant unless stated otherwise.

Child maltreatment mediates risk for poorer health and lower socioeconomic status. Child maltreatment partially mediated the association between growing up in a family affected by mental health problems and general health, life satisfaction and income in adulthood. The increased risk of more episodes of unemployment in adulthood was mediated fully by experiences of child maltreatment.

The association between growing up in a family affected by mental health problems or substance abuse and general health, life satisfaction, income and unemployment in adulthood is partly mediated by child maltreatment. Linear regression analyses showed no significant association between growing up in a family affected by mental health problems or substance abuse and BMI. Therefore, no mediation analysis was performed (see Table 4).

Discussion

This study examined long-term effects on health and socioeconomic status in adulthood related to growing up in a family affected by mental health problems or substance abuse in a sample that is representative for the German population in terms of age and gender. The results demonstrate increased risks for several socioeconomic, psychosocial and health-related problems of affected individuals. Furthermore, for the first time we were able to show a significant part of these effects are mediated by child maltreatment. These associations remained significant after adjustment for relevant sociodemographic confounders. These results suggest that the awareness of child maltreatment should be important issue in the treatment of adults with mental health problems or substance abuse.

The present analysis shows strongly increased risks for all forms of child maltreatment in families affected by mental health problems or substance abuse even after controlling for age, gender, and educational achievement as potential confounders. This is consistent with the results of other studies^{11,16,24–26}.

In addition, participants who grew up in a family affected by mental health problems or substance abuse, showed extensive health impairments in adulthood. While the increased risk for mental health problems is known²³, little literature exists on the consequences for physical health. However, in a study from the U.S. similar to this one, an increased risk for obesity was seen⁴.

While it was already known that children of mentally ill parents have a decreased quality of life²⁷ and poorer health²⁸, we were now able to show that the reduction in life satisfaction persists well into adulthood in a large representative sample. Nevertheless, the association of ACEs in general and reduced quality of life²⁹, reduced mental and physical health is widely known and well assessed^{2,30}—but these studies mainly assess the cumulative effect of ACEs or focus on child maltreatment¹, not assessing the effects of single household dysfunctions.

In our study, growing up in a family affected by mental health problems or substance abuses represents a considerable risk in various socio-economic areas: on average, participants stated more frequently that they had low levels of education and income. Interestingly, the latter even after controlling for level of education. This is consistent with results from earlier studies, which demonstrated a link between the mother's mental illness and lower school grades or later public transfer payments^{31,32}.

The associations between growing up in a family affected by mental health problems or substance abuse and long-term outcomes were all mediated by child maltreatment. Depending on the analysis, 29–41% of the strength of the relationships presented was mediated by child maltreatment—in the case of growing up in a family affected by mental health problems and later episodes of unemployment, the relationship was even completely mediated by child maltreatment.

In summary, this data suggests that in order for effective prevention of many long lasting negative effects to the live of affected children, the detection of and intervention to maltreatment in families affected by mental health problems or substance abuse are necessary. Therefore the issue of child maltreatment in the treatment of adult patients for mental health problems or substance abuse must be addressed comprehensively. This could be achieved by interdisciplinary approaches in healthcare institutions treating patients for mental health problems or substance abuse.

However, the mediation analysis also shows that the increased risk to health and socioeconomic status is only partly mediated by child maltreatment—there seem to be other factors that may also be relevant. Low socioeconomic status, social isolation and stigmatization due to mental health problems or substance abuse are both risk factors for, but also consequences of ACEs^{33,34} and thus forming a vicious circle. Another factor that may explain the increased risk for maltreatment demonstrated here is that in substance abusing parents, intoxication or withdrawal may result in impairments such as inconsistencies in the care and education of children and reduced impulse control³⁵. Neglect and impairment of consistent care³⁶, involvement of children in delusions and inappropriate affects that may cause insecurity in children³⁷, may be relevant in mentally ill parents. In addition, there is a risk of an impaired attachment to the child, which is of outstanding importance for the healthy development of children^{38,39}. Biological factors such as genetics in the intergenerational transmission of mental illness⁴⁰, altered health behavior such as smoking², and increased exposure to stress followed by changes in stress regulation⁴¹ may also play a role. The relevance of these factors can hardly be assessed in isolation and can vary depending on the background.

In families affected by mental health problems or substance abuse, these factors often accumulate. Although many parents are aware that they need support, they are often reluctant to accept professional help due to concerns about stigmatization or loss of custody⁴². Importantly, the risk of losing custody of children can be significantly reduced by early intervention⁴³—as can the risk of children developing mental disorders by themselves⁴⁴. A higher awareness of the existing risks for children of mentally ill or substance-abusing parents, combined with adequate support services, may thus contribute to the long-term destigmatization of affected parents through prevention.

Limitations. The assessment of ACEs was based on retrospective self-report, which can lead to underestimation of adverse childhood experiences due to recall bias, shame and misunderstandings^{45,46}. This in turn could lead to an underestimation of the results shown. In addition, it was not assessed which household member was mentally ill in the original household or who consumed substances, which substances were abused or which disease was present. Substance abuse and mental illness can be very heterogeneous. Additionally, severity of the individual types of maltreatment, the age at which they were experienced and for how long they were exposed to these experiences was not assessed. The wide age range of our sample is a strength. However, for some outcomes, such as unemployment, this may be a limitation as during the survey, some participants had not yet had the opportunity to be unemployed as they are still at school, for example. Even though missing data is scarce, only

valid cases were included into the analysis and selection bias can not be excluded. As this is an observational study, no causality can be derived.

Conclusion

Against the background of the here presented results, there is an urgent need for greater awareness for the increased risks of children of mentally ill and substance-abusing parents. Affected patients should be systematically asked about minor children and informed about the increased risks for affected children, just as about support offers. The here presented results indicate a need for better care for children of mentally ill and substance-abusing parents. Support for affected families, in which child protection plays a central role, and closer networking of psychiatrists, child and adolescent psychiatrists, paediatricians as well as youth welfare should be recommended.

Received: 30 January 2020; Accepted: 24 August 2020

Published online: 22 September 2020

References

1. Norman, R. E. *et al.* The long-term health consequences of child physical abuse, emotional abuse, and neglect: a systematic review and meta-analysis. *PLoS Med.* **9**, e1001349. <https://doi.org/10.1371/journal.pmed.1001349> (2012).
2. Felitti, V. J. *et al.* Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am. J. Prev. Med.* **14**, 245–258 (1998).
3. Clemens, V. *et al.* Association of child maltreatment subtypes and long-term physical health in a German representative sample. *Eur. J. Psychotraumatol.* **9**, 1510278. <https://doi.org/10.1080/20008198.2018.1510278> (2018).
4. Rehkopf, D. H. *et al.* Adverse childhood experiences and later life adult obesity and smoking in the United States. *Ann. Epidemiol.* **26**, 488–492.e485. <https://doi.org/10.1016/j.annepidem.2016.06.003> (2016).
5. Brown, D. W. *et al.* Adverse childhood experiences and the risk of premature mortality. *Am. J. Prev. Med.* **37**, 389–396. <https://doi.org/10.1016/j.amepre.2009.06.021> (2009).
6. Habetha, S., Bleich, S., Weidenhammer, J. & Fegert, J. M. A prevalence-based approach to societal costs occurring in consequence of child abuse and neglect. *Child Adolesc. Psychiatry Ment. Health* **6**, 35. <https://doi.org/10.1186/1753-2000-6-35> (2012).
7. Lee, R. T., Paulozzi, L. J., Melanson, C., Simon, T. R. & Arias, I. Child maltreatment surveillance: uniform definitions for public health and recommended data elements. Report. https://www.cdc.gov/violenceprevention/pdf/CM_Surveillance-a.pdf (Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, 2008).
8. Witt, A., Brown, R. C., Plener, P. L., Brähler, E. & Fegert, J. M. Child maltreatment in Germany: prevalence rates in the general population. *Child Adolesc. Psychiatry Ment. Health* **11**, 47. <https://doi.org/10.1186/s13034-017-0185-0> (2017).
9. Afzelius, M., Ostman, M., Rastam, M. & Priebe, G. Parents in adult psychiatric care and their children: a call for more interagency collaboration with social services and child and adolescent psychiatry. *Nord. J. Psychiatry* **72**, 31–38. <https://doi.org/10.1080/08039488.2017.1377287> (2018).
10. Wiegand-Greife, S., Matthejat, F. & Lenz, A. *Kinder psychisch kranker Eltern—Eine Einführung in die Thematik* (Vandenhoeck & Ruprecht, Göttingen, 2012).
11. Brockington, I. *et al.* WPA guidance on the protection and promotion of mental health in children of persons with severe mental disorders. *World Psychiatry* **10**, 93–102 (2011).
12. Brown, J., Cohen, P., Johnson, J. G. & Salzinger, S. A longitudinal analysis of risk factors for child maltreatment: findings of a 17-year prospective study of officially recorded and self-reported child abuse and neglect. *Child Abuse Negl.* **22**, 1065–1078. [https://doi.org/10.1016/S0145-2134\(98\)00087-8](https://doi.org/10.1016/S0145-2134(98)00087-8) (1998).
13. Mulder, T. M., Kuiper, K. C., van der Put, C. E., Stams, G. J. M. & Assink, M. Risk factors for child neglect: a meta-analytic review. *Child Abuse Negl.* **77**, 198–210. <https://doi.org/10.1016/j.chiabu.2018.01.006> (2018).
14. Felitti, V. J. *et al.* Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the adverse childhood experiences (ACE) study. *Am. J. Prev. Med.* **14**, 245–258. [https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8) (1998).
15. Gilbert, R. *et al.* Burden and consequences of child maltreatment in high-income countries. *The Lancet* **373**, 68–81. [https://doi.org/10.1016/S0140-6736\(08\)61706-7](https://doi.org/10.1016/S0140-6736(08)61706-7) (2009).
16. Clemens, V. *et al.* Child maltreatment is mediating long-term consequences of household dysfunction in a population representative sample. *Eur. Psychiatry* **58**, 10–18. <https://doi.org/10.1016/j.eurpsy.2019.01.018> (2019).
17. Clemens, V. *et al.* Authoritarianism and the transgenerational transmission of corporal punishment. *Child Abuse Negl.* **106**, 104537. <https://doi.org/10.1016/j.chiabu.2020.104537> (2020).
18. Wingenfeld, K. *et al.* Reliable, valide und ökonomische Erfassung früher Traumatisierung: Erste psychometrische Charakterisierung der deutschen Version des Adverse Childhood Experiences Questionnaire (ACE). *Psychother. Psychosom. Med.* **61**, e10–e14. <https://doi.org/10.1055/s-0030-1263161> (2011).
19. Beierlein, C., Kovaleva, A., László, Z., Kemper, C. J., & Rammstedt, B. *Kurzskala zur Erfassung der Allgemeinen Lebenszufriedenheit (L-1). Zusammenstellung sozialwissenschaftlicher Items und Skalen*, 2015.
20. EuroQol Group. EuroQol—a new facility for the measurement of health-related quality of life. *Health Policy* **16**, 199–208 (1990).
21. Oppe, M. V. R. M. *EQ-5D-3L User Guide. Basic Information on How to Use the EQ-5D-3L Instrument*, 2015.
22. Hinze, A., Klaiberg, A., Brähler, E. & König, H. H. The Quality of Life Questionnaire EQ-5D: modelling and norm values for the general population. *Psychother. Psychosom. Med. Psychol.* **56**, 42–48. <https://doi.org/10.1055/s-2005-867061> (2006).
23. Hayes, A. F. *Introduction to Mediation, Moderation, and Conditional Process Analysis A Regression-Based Approach*. (Guilford Publications, Inc., New York, 2013).
24. C. D. in *Kindesmisshandlung und Vernachlässigung. Ein Handbuch*. 141–154 (Deegener G und Körner W 2005).
25. Dubowitz, H. *et al.* Identifying children at high risk for a child maltreatment report. *Child Abuse Negl.* **35**, 96–104. <https://doi.org/10.1016/j.chiabu.2010.09.003> (2011).
26. Dong, M. *et al.* The interrelatedness of multiple forms of childhood abuse, neglect, and household dysfunction. *Child Abuse Negl.* **28**, 771–784. <https://doi.org/10.1016/j.chiabu.2004.01.008> (2004).
27. Ravens-Sieberer, U., Ellert, U. & Erhart, M. Gesundheitsbezogene Lebensqualität von Kindern und Jugendlichen in Deutschland: Eine Normstichprobe für Deutschland aus dem Kinder- und Jugendgesundheitsurvey (KIGGS). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. <https://doi.org/10.1007/s00103-007-0244-4> (2007).
28. Hagen, K. A., Hilsen, M., Kallander, E. K. & Ruud, T. Health-related quality of life (HRQoL) in children of ill or substance abusing parents: examining factor structure and sub-group differences. *Qual. Life Res.* <https://doi.org/10.1007/s11136-018-2067-1> (2018).
29. Salinas-Miranda, A. A. *et al.* Adverse childhood experiences and health-related quality of life in adulthood: revelations from a community needs assessment. *Health Qual. Life Outcomes* **13**, 123. <https://doi.org/10.1186/s12955-015-0323-4> (2015).

30. Holman, D. M. *et al.* The association between adverse childhood experiences and risk of cancer in adulthood: a systematic review of the literature. *Pediatrics* **138**, S81–S91. <https://doi.org/10.1542/peds.2015-4268L> (2016).
31. Netsi, E. *et al.* Association of persistent and severe postnatal depression with child outcomes. *JAMA Psychiatry* **75**, 247–253. <https://doi.org/10.1001/jamapsychiatry.2017.4363> (2018).
32. Homlong, L., Rosvold, E. O., Sagatun, A., Wentzel-Larsen, T. & Haavet, O. R. Living with mentally ill parents during adolescence: a risk factor for future welfare dependence? A longitudinal, population-based study. *BMC Public Health* **15**, 413. <https://doi.org/10.1186/s12889-015-1734-1> (2015).
33. Goodman, S. H. Depression in mothers. *Annu. Rev. Clin. Psychol.* **3**, 107–135. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091401> (2007).
34. van Santvoort, F. *et al.* The impact of various parental mental disorders on children's diagnoses: a systematic review. *Clin. Child Fam. Psychol. Rev.* **18**, 281–299. <https://doi.org/10.1007/s10567-015-0191-9> (2015).
35. Beck, A. & Heinz, A. Alcohol-related aggression—social and neurobiological factors. *Deutsches Ärzteblatt International* **110**, 711–715. <https://doi.org/10.3238/arztebl.2013.0711> (2013).
36. Riley, A. W. *et al.* Development of a family-based program to reduce risk and promote resilience among families affected by maternal depression: theoretical basis and program description. *Clin. Child Fam. Psychol. Rev.* **11**, 12–29. <https://doi.org/10.1007/s10567-008-0030-3> (2008).
37. Bosanac, P., Buist, A. & Burrows, G. Motherhood and schizophrenic illnesses: a review of the literature. *Aust. N. Z. J. Psychiatry* **37**, 24–30. <https://doi.org/10.1046/j.1440-1614.2003.01104.x> (2003).
38. Parolin, M. & Simonelli, A. Attachment theory and maternal drug addiction: the contribution to parenting interventions. *Front. Psychiatry* **7**, 152. <https://doi.org/10.3389/fpsy.2016.00152> (2016).
39. Widom, C. S., Czaja, S. J., Kozakowski, S. S. & Chauhan, P. Does adult attachment style mediate the relationship between childhood maltreatment and mental and physical health outcomes?. *Child Abuse Negl.* **76**, 533–545. <https://doi.org/10.1016/j.chiabu.2017.05.002> (2018).
40. Rasic, D., Hajek, T., Alda, M. & Uher, R. Risk of mental illness in offspring of parents with schizophrenia, bipolar disorder, and major depressive disorder: a meta-analysis of family high-risk studies. *Schizophr. Bull.* **40**, 28–38. <https://doi.org/10.1093/schbul/sbt114> (2014).
41. Carpenter, L. L. *et al.* Decreased adrenocorticotropic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biol. Psychiatry* **62**, 1080–1087. <https://doi.org/10.1016/j.biopsych.2007.05.002> (2007).
42. Jungbauer, J., Stelling, K., Kuhn, J. & Lenz, A. How do mothers and fathers suffering from schizophrenia experience their parenthood? Results from an in-depth interview study. *Psychiatr. Prax.* **37**, 233–239. <https://doi.org/10.1055/s-0029-1223535> (2010).
43. Seeman, M. V. Intervention to prevent child custody loss in mothers with schizophrenia. *Schizophr. Res. Treat.* **2012**, 6. <https://doi.org/10.1155/2012/796763> (2012).
44. Siegenthaler, E., Munder, T. & Egger, M. Effect of preventive interventions in mentally ill parents on the mental health of the offspring: systematic review and meta-analysis. *J. Am. Acad. Child Adolesc. Psychiatry* **51**, 8–17.e18. <https://doi.org/10.1016/j.jaac.2011.10.018> (2012).
45. Gilbert, R. *et al.* Burden and consequences of child maltreatment in high-income countries. *Lancet* **373**, 68–81. [https://doi.org/10.1016/s0140-6736\(08\)61706-7](https://doi.org/10.1016/s0140-6736(08)61706-7) (2009).
46. Fergusson, D. M., Horwood, L. J. & Woodward, L. J. The stability of child abuse reports: a longitudinal study of the reporting behaviour of young adults. *Psychol. Med.* **30**, 529–544. <https://doi.org/10.1017/s0033291799002111> (2000).

Acknowledgements

We thank all participants and staff involved in data retrieval and management.

Author contributions

All authors have made substantial contributions to this work and have read and approved the final manuscript as submitted. They agree to be accountable for all aspects of the work. All authors were involved in conceiving and designing the study and interpreting the data. In Detail, V.C. analyzed the data. V.C., O.B., A.W. and C.S. drafted the article. J.M.F. and P.L.P. designed the research idea. E.B. and B.S. supervised the project and data collection.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit-sectors. VC, OB, CS, AW, EB and BS state that they have no conflict of interests. JMF has received research funding from the EU, DFG (German Research Foundation), BMG (Federal Ministry of Health), BMBF (Federal Ministry of Education and Research), BMFSFJ (Federal Ministry of Family, Senior Citizens, Women and Youth), BfArM (Federal Institute for Drugs and Medical Devices), German armed forces, several state ministries of social affairs, State Foundation Baden-Württemberg, Volkswagen Foundation, Pontifical Gregorian University, CJD, Caritas, Diocese of Rottenburg-Stuttgart. Moreover, he received travel grants, honoraria and sponsoring for conferences and medical educational purposes from DFG, AACAP, NIMH/NIH, EU, Pro Helvetia, Shire, several universities, professional associations, political foundations, and German federal and state ministries during the last 5 years. Every grant and every honorarium has to be declared to the law office of the University Hospital Ulm. Professor Fegert holds no stocks of pharmaceutical companies. PLP has received research funding from the German Federal Institute for Drugs and Medical Devices (BfArM), German Federal Ministry of Education and Research (BMBF), VW-Foundation, Baden-Württemberg Foundation, Lundbeck, Servier. He received a speaker's honorarium from Shire. Professor Plener holds no stocks of pharmaceutical companies. Open Access funding provided by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to V.C.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020